


Fig. 5. Hall resistance for low hole concentration (5×10^{10} to $8 \times 10^{10} \text{ cm}^{-2}$) in a tetracene MIS device at 1.7 K. Distinct integer Hall plateaus and the fractional quantum Hall states $j = 1/3$, $j = 2/3$, and $j = 2/5$ are clearly visible.



samples with mobilities of several million square centimeters per volt per second. In addition, the combination of a reduced dielectric constant and a much higher effective mass than in GaAs makes new parameter regions accessible, enabling studies of the physics of strongly interacting electron systems, such as the metal-insulator transition in two dimensions (19).

We have presented ample experimental evidence for band-like charge transport in delocalized states in these organic semiconductors: the high mobilities observed at low temperatures, the temperature dependence of the mobility, the large effective bandwidth at low temperatures ($W \gg k_B T$, where k_B is the Boltzmann constant), Shubnikov–de Haas oscillations, and the observation of the integer and fractional QHE. All these observations of transport properties, particularly at low temperatures, are similar to those observed in conventional inorganic semiconductors. It appears that the measured effective masses of order 1 to 1.5 m_e are not those of bare holes or electrons, but instead are charge carriers dressed by a polarization cloud. Looking forward, then, one might expect the adoption of additional concepts from inorganic semiconductor technology (such as superlattices and quantum wells) to yield interesting electronic and optical properties. Layered organic semiconductors (17, 19, 20) could offer more latitude for device engineering because they are van der Waals–bonded, and hence minimal constraints are imposed by lattice matching (21, 22). Furthermore, interesting new phenomena can be anticipated to result from stronger electron-phonon interaction or strong transport anisotropy, opening up a new field of research.

References and Notes

1. K. von Klitzing, G. Dorda, M. Pepper, *Phys. Rev. Lett.* **45**, 494 (1980).
2. H. L. Stormer, *Rev. Mod. Phys.* **71**, 875 (1999); D. C. Tsui, H. L. Stormer, A. C. Gossard, *Phys. Rev. B* **25**, 1408 (1982).
3. M. Ribault *et al.*, *J. Phys. Lett.* **45**, L935 (1984).
4. R. V. Chamberlin *et al.*, *Phys. Rev. Lett.* **60**, 1189 (1988).
5. P. M. Chaikin, W. Kang, S. Hannahs, R. C. Yu, *Physica B* **177**, 353 (1992).

6. For another organic system, α -(BEDT-TTF)₂MHg(SCN)₄ salts [where BEDT-TTF is bis(ethylenedithio)-tetrathiafulvalene and M = K or Tl], indirect evidence of a QHE was found in high-field de Haas–van Alphen measurements; this was supported by further evidence from Hall transport measurements. These phenomena in charge-transfer salts, however, are ascribed to a bulk QHE substantially different from the “conventional” QHE in a 2D electron gas in an inorganic heterostructure or in a MIS structure [N. Harrison *et al.*, *Phys. Rev. Lett.* **77**, 1576 (1996); M. M. Honold *et al.*, *Phys. Rev. B* **59**, R10417 (1999)].

7. Ch. Kloc, P. G. Simpkins, T. Siegrist, R. A. Laudise, *J. Cryst. Growth* **182**, 416 (1997).
8. R. A. Laudise, Ch. Kloc, P. G. Simpkins, T. Siegrist, *J. Cryst. Growth* **187**, 449 (1998).
9. J. H. Schön, S. Berg, Ch. Kloc, B. Batlogg, *Science* **287**, 1022 (2000).
10. A. Dodabalapur, L. Torsi, H. E. Katz, *Science* **268**, 270 (1995).
11. N. Karl, J. Marktanner, R. Stehle, W. Warta, *Synth. Met.* **41–43**, 2473 (1991).
12. D. M. Burland and U. Konzelmann, *J. Chem. Phys.* **67**, 319 (1977).
13. K. von Klitzing, *Rev. Mod. Phys.* **58**, 519 (1986).
14. P. M. Littlewood, personal communication.
15. J. Cornil, personal communication.
16. W. Warta and N. Karl, *Phys. Rev. B* **32**, 1172 (1985).
17. D. C. Singh and S. C. Matur, *Mol. Cryst. Liq. Cryst.* **27**, 55 (1974).
18. R. C. Haddon, personal communication.
19. S. V. Kravchenko, G. V. Kravchenko, J. E. Furneaux, V. M. Pudalov, M. D’Iorio, *Phys. Rev. B* **50**, 8039 (1994).
20. H. Akimichi, T. Inoshita, S. Hotta, H. Noge, H. Sakaki, *Appl. Phys. Lett.* **63**, 3158 (1993).
21. T. Minakata and Y. Mori, *Pol. Adv. Technol.* **6**, 611 (1995).
22. F. F. So and S. R. Forrest, *Phys. Rev. Lett.* **66**, 2649 (1991).
23. We thank E. Bucher for use of his equipment, and A. Dodabalapur, J. W. P. Hsu, H. Y. Hwang, H. L. Störmer, and C. M. Varma for helpful discussions. Supported by the Deutsche Forschungsgemeinschaft (J.H.S.).

6 April 2000; accepted 19 May 2000

Discovery of a High-Energy Gamma-Ray-Emitting Persistent Microquasar

Josep M. Paredes,^{1*} Josep Martí,² Marc Ribó,¹ Maria Massi³

Microquasars are stellar x-ray binaries that behave as a scaled-down version of extragalactic quasars. The star LS 5039 is a new microquasar system with apparent persistent ejection of relativistic plasma at a 3-kiloparsec distance from the sun. It may also be associated with a γ -ray source discovered by the Energetic Gamma Ray Experiment Telescope (EGRET) on board the COMPTON–Gamma Ray Observatory satellite. Before the discovery of LS 5039, merely a handful of microquasars had been identified in the Galaxy, and none of them was detected in high-energy γ -rays.

The $V = 11.2$ magnitude star LS 5039 (1) has been recently identified as a nearby high-mass x-ray binary with spectral type O7V(f) (2) and persistent radio emission (3, 4). Here, we report high-resolution radio observations with the Very Long Baseline Array (VLBA) and the Very Large Array (VLA) that reveal that LS 5039 is resolved into bipolar radio jets emanating from a central core.

Because LS 5039 appeared unresolved ($\leq 0.1''$) to the VLA alone, we proceeded to study this object with milliarc sec resolution using the VLBA at the frequency of 5 GHz (6-cm wavelength) on 8 May 1999. The VLA in its phased array mode, equivalent to a dish of 115-m diameter, also participated as an independent station, providing sensitive baselines with the VLBA antennas. The source 3C345 was used as a fringe-finder, whereas J1733–1304 was the phasing source for the VLA. The data were calibrated using standard procedures in unconnected radio interferometry. The resulting pattern of the observed visibility amplitudes, decaying as a function of baseline length, indicated that LS 5039 had structure at milliarc sec scales.

The final synthesis map (Fig. 1) shows that bipolar jets emerge from a central core. A de-

¹Departament d’Astronomia i Meteorologia, Universitat de Barcelona, Av. Diagonal 647, E-08028 Barcelona, Spain. ²Departamento de Física, Escuela Politécnica Superior, Universidad de Jaén, Calle Virgen de la Cabeza 2, E-23071 Jaén, Spain. ³Max Planck Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany.

*To whom correspondence should be addressed. E-mail: josemp@am.ub.es

REPORTS

convolved angular size of about 2 milliarc sec is estimated for the core. The jets extend over 6 milliarc sec on the sky oriented along a position angle (PA) of 125° with respect to the north, and they account for 20% of the total 16-mJy flux density. To obtain some order of magnitude estimates, we will assume that the overall size of the radio source is approximately 6×2 milliarc sec². This implies a high brightness temperature of $\sim 9.4 \times 10^7$ K, indicating synchrotron radiation. The LS 5039 radio spectrum as a function of frequency ν , namely $S_\nu \propto \nu^\alpha$, often displays a negative spectral index $\alpha = -0.5$ in agreement with a nonthermal optically thin emission mechanism (3, 4). The detection of jets occurred at a time when the source was at its typical persistent level of radio emission, and only moderately variable, as inferred from concurrent radio monitoring by the Green Bank Interferometer (GBI) (Fig. 2). The absence of any precursor outburst for the radio jets strongly suggests that they are always present and continuously emanating from the core. The flux density ratio between the southeast and northwest jet components is estimated as 2.1 ± 0.4 . It seems reasonable that this brightness asymmetry reflects a relativistic Doppler boosting effect (5). If a continuous jet flow is assumed, the projected velocity required is then $v \cos \theta = (0.15 \pm 0.04)c$, where c is the speed of light and θ the ejection angle with the line of sight. It is straightforward to then derive a lower and upper limit for the jet velocity [$v \geq (0.15 \pm 0.04)c$] and the ejection angle ($\theta \leq 81^\circ \pm 2^\circ$), respectively.

X-ray binaries with collimated radio jets belong to the class of galactic microquasars. The production of jets is almost certainly related to the capture of matter from a normal star by a black hole or neutron star companion. This is a highly energetic process with observable consequences from radio to hard x-rays (6) and possibly beyond. The recent third EGRET catalog of high-energy ($E_\gamma > 100$ MeV) γ -ray sources (7) contains nearly 100 unidentified emitters at low galactic latitudes. The position of LS 5039 is well inside the 95% confidence contour of the EGRET source 3EG J1824-1514, whose radius is about 0.5° . Moreover, LS 5039 is the only x-ray emitter within 1° of 3EG J1824-1514 listed in the ROSAT (Roentgen Satellite) All Sky Survey (8). Such a good position agreement between an EGRET source and a peculiar radio jet x-ray binary strongly implies that both objects are the same. Thus, this microquasar system is likely associated with an EGRET source. The γ -ray emission observed reveals a rather persistent flux of >100 MeV photons for the last 10 years (Fig. 3).

Using modern photometric data (9) and the reddening free parameter formulation (10), we obtained a distance estimate of 3.1 kpc. This value is in excellent agreement with

independent results based on the star color excess (2). On the other hand, a common intrinsic radio luminosity has been recently suggested for persistent x-ray binaries (11). According to this, the LS 5039 average flux density of a few tens of mJy at cm wavelengths would imply a rough distance value not higher than 2 kpc. Thus, different distance indicators show that LS 5039 is nearby, and we adopt a distance of 3 kpc. Therefore, this star appears to be one of the closest and optically brightest microquasars among the persistent members of this class. Several other nontransient microquasars happen to be beyond distances of about 8 kpc (12), such as the prototypical 1E 1740.7-2942 in the heavily obscured regions of the Galactic Center (13).

The synchrotron radio luminosity between 0.1 and 100 GHz for this distance is $L_{\text{rad}} \sim 7.5 \times 10^{30}$ erg s⁻¹. The average γ -ray flux for all EGRET viewing periods in Fig. 3 is $\Phi_\gamma = (35.2 \pm 6.5) \times 10^{-8}$ photon cm⁻² s⁻¹ with photon spectral index $p = 2.19 \pm 0.18$, where $\Phi_\gamma \propto E_\gamma^{-p}$. The EGRET photon index of LS 5039 is practically identical to that of

1E 1740.7-2942, i.e., steeper than the $p < 2$ values usually found for pulsars (14). The corresponding integrated luminosity amounts to $L_\gamma (>100 \text{ MeV}) \sim 3.8 \times 10^{35}$ erg s⁻¹, compared to an x-ray luminosity (4) of $L_x(1.5 - 12 \text{ keV}) \sim 5 \times 10^{34}$ erg s⁻¹. Additional information on the source energetics can be obtained by assuming energy equipartition between the relativistic electrons and the magnetic field (15). We are forced to use the overall source parameters observed because not enough information is yet available for appropriate calculations in the rest frame of the ejecta. The corresponding results are nevertheless expected to be within an order of magnitude for a mildly relativistic system. Under these assumptions, the observed radio properties of LS 5039 imply a total energy content in relativistic electrons of $E_e \sim 4.8 \times 10^{39}$ erg, with an equipartition magnetic field of ~ 0.2 G.

While flowing away into opposite jets, the relativistic electrons are exposed to a huge output of ultraviolet (UV) photons from the hot optical star. Thus, it appears likely that a significant fraction of the EGRET emission




Fig. 1. High-resolution radio map of the nearby star LS 5039 obtained with the VLBA and the VLA in phased array mode at 6-cm wavelength. The presence of radio jets in this high-mass x-ray binary is the main evidence supporting its microquasar nature. The contours shown correspond to 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 40, and 50 times 0.085 mJy per beam, the rms noise. The ellipse at the bottom right corner represents the half-power beam width of the synthesized beam, 3.4×1.2 (milliarc sec²) with a PA of 0° . The map is centered at the LS 5039 position $\alpha_{J2000} = 18^{\text{h}}26^{\text{m}}15.056^{\text{s}}$ and $\delta_{J2000} = -14^\circ50'54.24''$. North is at the top and east is at the left. One milliarc sec is equivalent to 4.5×10^{13} cm (3 AU) for a distance of 3 kpc.




Fig. 2. GBI radio monitoring of LS 5039, at 2.25 GHz (13 cm), during the weeks before and after the date of our VLBA+VLA observation, indicated by the vertical bar. No strong flaring event was recorded, suggesting that the presence of radio jets must be a permanent feature of LS 5039.

arises as a result of inverse Compton (IC) scattering of these photons by the same radio-emitting electrons. The energy shift in this process is such that $E_\gamma \sim \gamma_e^2 E_{\text{ph}}$, where the energies of the γ -ray and the stellar photon are related through the squared Lorentz factor of the relativistic electron. For an O7 main sequence star, a UV luminosity of $L_* \sim 10^{38}$ erg s^{-1} is expected to be mostly emitted by photons with $E_{\text{ph}} \sim 10$ eV. In order to scatter them into γ -rays with $E_\gamma \sim 100$ MeV, electrons with Lorentz factors $\sim 10^3$, equivalent to energies of $\sim 10^{-3}$ erg, are needed. Considering the persistent EGRET luminosity, the lifetime of such electrons against dominant IC losses will be $t_c \sim E_e/L_\gamma \sim 1.3 \times 10^4$ s.

The electron energy will decay with time by IC scattering according to (15) (in centimeter-gram-second units)

$$\left(\frac{dE}{dt}\right)_{\text{IC}} = 3.97 \times 10^{-2} U_{\text{rad}} E^2 \quad (1)$$

where U_{rad} is the UV radiation energy density. For electrons flowing away into jets, assumed perpendicular to the plane of a circular orbit with radius r , we have $U_{\text{rad}} = L_*/4\pi c(r^2 + v^2 t^2)$ at a time t after injection. For an electron with initial energy E_0 , its IC lifetime can be expressed as $t_c = 25.2/U_{\text{rad}} E_0$ when injected into the jet basis close to the compact object. This implies then that the γ -ray-emitting electrons must be initially exposed to $U_{\text{rad}} \sim 2.0$ erg cm^{-3} . Such values of radiation energy density are available if the jets originate at a distance $r \sim 1.2 \times 10^{13}$ cm from the star.

Equation 1 can be solved to give

$$E(t) = \frac{E_0}{1 + (r/vt_c) \arctan(vt/r)} \quad (2)$$

By imposing the condition that electrons with $E_0 \sim 10^{-3}$ erg are able to abandon the region of heavy IC emission in the star vicinity, the condition $\pi r/2vt_c < 1$ must be fulfilled so that they still retain enough energy to power

the extended radio jets. This requirement allows us to constrain the jet velocity to values $v > 0.05c$, in agreement with the previous discussion of Doppler boosting. The true jet velocity is not likely to exceed a mildly relativistic value $v \sim 0.4c$, which we crudely estimate assuming that the 6 milliarc sec extended jets have to be replenished in a t_c time. The Lorentz factor of the jets would then be $\gamma_v = 1/\sqrt{1 - (v/c)^2} \sim 1.1$, i.e., not extremely relativistic. The size of the region where γ -rays are produced in this scenario is $vt_c > 1.8 \times 10^{13}$ cm, i.e., larger than the orbital radius.

The central engine in LS 5039 must be supplying $\dot{E}_e \sim L_\gamma \sim 3.8 \times 10^{35}$ erg s^{-1} in the form of relativistic electrons. Their energy distribution is expected to be a power law $kE^{2\alpha-1} dE = kE^{-2} dE$ to produce the observed spectral index. Assuming electron energies in the range $m_e c^2 \leq E \leq \gamma_{\text{max}} m_e c^2$ we have $\dot{E}_e = k \int E^{2\alpha-1} dE = k \ln \gamma_{\text{max}}$. Therefore, if the proton mass is $m_p \approx 1800 m_e$ for every relativistic electron, the proton mass flow into the jets can be written as $\dot{M}_{\text{jet}} = m_p k \int E^{2\alpha-1} dE \approx 1800 \dot{E}_e / c^2 \ln \gamma_{\text{max}} \sim 1.3 \times 10^{-9} M_\odot \text{ year}^{-1}$. The equivalent kinetic luminosity is $L_K = (\gamma_v - 1) \dot{M}_{\text{jet}} c^2 \sim 10^{37}$ erg s^{-1} , which is weakly dependent on the maximum energy cutoff assumed ($\gamma_{\text{max}} \sim 10^4$). This kinetic power is about four orders of magnitude less than that estimated for the strong ejections of the superluminal microquasar GRS 1915+105 (16). Both the mass outflow and kinetic energy estimates would not be significantly affected if positrons are considered instead of protons because the relativistic mass of an electron with Lorentz factor $\sim 10^3$ is comparable to m_p .


LS 5039 is one of the nearest microquasars to be discovered. It has strong high-energy γ -ray emission, which sets limits on the likely velocity of its jets via IC energy losses. Most of known microquasars were discovered only after undergoing a noticeable outburst that triggered detection by the bat-

tery of satellites and ground-based observatories. Some recent examples include CI Camelopardalis (17) and the nearby transient V4641 Sagittarii (18). The microquasar nature of these two objects is tantalizing in that both are bright optical stars. CI Camelopardalis was even cataloged as a variable star before its outburst. Therefore, a careful examination of modern archive databases may reveal a previously unnoticed population of microquasars. Indeed, our identification of LS 5039 as a potential candidate resulted from a systematic cross-correlation between public archives of astrophysical data in the x-ray, radio, and optical domains (8, 19, 20). The success of this approach for systematic identification opens the possibility of new findings which may confirm that the microquasar phenomenon is not as rare as it seems.

References and Notes

1. C. B. Stephenson and N. Sanduleak, *Publ. Warner Swasey Obs.* **1**, 1 (1971).
2. C. Motch, F. Haberl, K. Dennerl, M. Pakull, E. Janot-Pacheco, *Astron. Astrophys.* **323**, 853 (1997).
3. J. Martí, J. M. Paredes, M. Ribó, *Astron. Astrophys.* **338**, L71 (1998).
4. M. Ribó, P. Reig, J. Martí, J. M. Paredes, *Astron. Astrophys.* **347**, 518 (1999).
5. T. J. Pearson and J. A. Zensus, in *Superluminal Radio Sources*, J. A. Zensus and T. J. Pearson, Eds. (Cambridge Univ. Press, Cambridge, 1987), pp. 1–11.
6. I. F. Mirabel and L. F. Rodríguez, *Annu. Rev. Astron. Astrophys.* **39**, 409 (1999).
7. R. C. Hartman et al., *Astrophys. J. Suppl. Ser.* **123**, 79 (1999).
8. W. Voges et al., *Astron. Astrophys.* **349**, 389 (1999).
9. J. F. LaHulla and J. Hilton, *Astron. Astrophys. Suppl. Ser.* **94**, 265 (1992).
10. B. C. Reed, *Publ. Astron. Soc. Pac.* **105**, 1465 (1993).
11. R. P. Fender and M. A. Hendry, *Mon. Not. R. Astron. Soc.*, in press.
12. J. Greiner, in *Cosmic Explosions*, Proceedings of the 10th Annual Astrophysics Conference in Maryland, College Park, MD, 11 to 13 October 1999, S. Holt and W. W. Zhang, Eds., in press.
13. I. F. Mirabel, L. F. Rodríguez, B. Cordier, J. Paul, F. Lebrun, *Nature* **358**, 215 (1992).
14. M. Merk et al., *Astron. Astrophys. Suppl. Ser.* **120**, 465 (1996).
15. A. G. Pacholczyk, *Radio Astrophysics* (Freeman, San Francisco, CA, 1970).
16. I. F. Mirabel and L. F. Rodríguez, *Nature* **371**, 46 (1994).
17. R. M. Hjellming and A. M. Mioduszewski, *Int. Astron. Union Circ.* **6872** (1998).
18. R. M. Hjellming et al., *Int. Astron. Union Circ.* **7265** (1999).
19. J. J. Condon et al., *Astron. J.* **115**, 1693 (1998).
20. B. M. Lasker et al., *Astron. J.* **99**, 2019 (1990).
21. D. J. Helfand, S. Zoonematkermani, R. H. Becker, R. L. White, *Astrophys. J. Suppl. Ser.* **80**, 211 (1992).
22. This research was partly supported by the Dirección General de Enseñanza Superior e Investigación Científica (PB97-0903) in Spain. M.R. acknowledges receipt of a fellowship from Generalitat de Catalunya. M.M. and J.M. also acknowledge support by the European Commission's Training and Mobility of Researchers program and Junta de Andalucía, respectively. The VLBA and VLA are operated by the National Radio Astronomy Observatory (NRAO) with Associated Universities Inc. and are funded by the NSF. The GBI is a facility of the NSF operated by NRAO with support from the NASA High-Energy Astrophysics program.

Fig. 3. Radio and γ -ray light curves of LS 5039 and 3EG J1824-1514, which we propose originate in the same object. Both LS 5039 and 3EG J1824-1514 are consistent with a persistent level of emission over the last decade. The fluxes plotted here are taken from the literature and archive data (3, 4, 19, 27). Error bars for GBI (± 4 mJy) are not shown for clarity, whereas those of the VLA are usually smaller than the symbol size.



29 March 2000; accepted 15 May 2000